

# DonJoy<sup>®</sup> ISOFORM<sup>™</sup> LO

Clinician Application Instructions

BEFORE USING THE DEVICE, PLEASE READ THE INSTRUCTIONS FOR USE COMPLETELY AND CAREFULLY. CORRECT APPLICATION IS VITAL TO THE PROPER FUNCTIONING OF THE DEVICE.

# SIZING:

## Step 1. Size the Brace

Begin by measuring the circumference of the patient approximately 1" (2.5cm) above the widest part of the hips. Then, match the patient's size to the sizing chart below. If patient's circumferential measurement is between sizes, select the larger brace.

SIZE	CIRCUMFERENTIAL MEASUREMENT
S/M	Less than 48" (122 cm)
L/XL	49" - 60" (125 cm - 152 cm)

**NOTE:** It is not uncommon for the patient to experience post-surgical swelling, followed by weight loss (10-20+ lbs/4.5-9.1 kg). The LO+ brace is designed with two (2) adjustable belts which can be easily sized to achieve optimal fit.

# FITTING:

### Adjust Circumference

To adjust waist belt circumference, align the tip of the hook and loop attachment on the rear panel section with the correct circumference. Repeat for opposite side. (Fig. A) For patients whose circumferential measurement is between 60" and 70", (152cm and 178cm), use optional bridge wing (sold separately). Apply hook portion of donning mitt to the loop exterior.

# **APPLICATION:**

### Step 2. Loosen the Brace

Detach the pull tabs from the loop exterior. Pull the brace in opposite directions. (Fig. B)

### Step 3. Position the Brace

Center the rear section of the belt at the midline of the lower back, so that the belt is just above the crease of the buttocks. Note: The brace is intended to be worn over clothing.

### Step 4. Secure the Brace

Secure the hook portion of the donning mitt to the loop exterior. (Fig. C) Before tightening the brace, make sure the rear panel section is positioned evenly on both sides of the patient. Have the patient slip their thumbs through the holes in the pull tabs and pull the tabs away from the body. **NOTE FOR OPTIMAL WAIST FIT:** The belt wings may be secured to the rear panel section at an angle to accommodate a size differential between the waist and hips. (Fig. D) When tightened, secure the pull tabs to the loop exterior. Adjust compression as needed. (Fig. E)

### Helpful Tip: Brace Migration

To reduce brace migration, follow these easy steps. Before sitting, loosen the brace by detaching the pull tabs. Once seated, tighten the pull tabs by pulling across the body and re-adhering to the loop exterior. When standing up, readjust pull tabs.


### Brace Removal

To remove the brace, **first** loosen it by detaching the pull tab from the loop exterior. Then unhook the donning mitt from the loop exterior and take off the brace.


### **Brace Re-application**


When reapplying the brace, it is easiest to guide the belt by inserting a hand in the donning mitt pocket.















# DonJoy<sup>®</sup> ISOFORM<sup>™</sup> LO

Clinician Application Instructions

## HOT/COLD PACK (sold separately):

If patient will use a hot/cold pack with the LO+, follow the instructions on the hot/cold pack. Insert hot/cold pack into pocket and adhere to back panel.

## ADJUSTING PULL TAB LENGTH (OPTIONAL)

Remove lace spool from the pull tab pocket and wind lace around the spool to shorten the length of the pull tab. Unwind lace from the spool to lengthen the pull tab. (Figure F) Insert lace spool back into pull tab pocket.

## ATTACH/REMOVE BACK PANEL (FOR CLEANING)

Open tall rear panel pouch from waist belt. Detach rear panel section by unlooping waist belt. Remove panel. (Figure G) Reattach rear panel section by looping waist belt through panel. With tall rear panel pouch open, lay waist belt on top of lacing guides. Close pouch.

## **CLEANING INSTRUCTIONS:**

Separate the brace's removable REAR PANEL before cleaning. Hand wash in cold water, less than 86°F (30°C), using mild soap. Air dry.

NOTE: If not rinsed thoroughly, residual soap may cause irritation and deteriorate material.

Do NOT iron. Do NOT tumble dry. Do NOT bleach.

### INTENDED USER PROFILE:

The user should be able to: Read and understand the directions, warnings and cautions.

### INTENDED USE/INDICATIONS:

Acute and chronic low back pain, sciatica, lower back sprains or strains, lumbar disc displacement, osteoporosis, disc herniation and degeneration, post-operative laminectomy, post-operative discectomy, spondylolisthesis, pre and post surgical care.

**CONTRAINDICATIONS:** Pregnancy and circulation, pulmonary, cardiovascular or skeletal conditions which have risk to be made worse as a result of compression and/or pressure.

WARNINGS AND CAUTIONS: If you experience any pain, swelling, sensation changes, excessive discoloration, irritation to the skin or any unusual reaction while using this product, consult your medical professional. If any distortion of the support panels, fraying or deterioration of lace cables is noticed, discontinue use. Brace is intended to be worn over clothing. Brace should be appropriately sized to the patient to ensure proper support and function. If the brace does not fit properly, it may be reshaped and adjusted by a healthcare professional. Care should be taken with patients, especially the elderly, not to apply the brace too tightly, which MAY cause excessive pressure. Patients should be instructed on appropriate tensioning during ongoing use. Patient should see a healthcare professional for a replacement if a product quality issue occurs within the warranty timeframe. Brace should not be exposed to moisturizers, chemicals, or solvents, as they may affect durability.

### MATERIAL CONTENTS:

Nylon, Polyester, Polyethylene, Polyurethane, Cotton, Acrylic, Polypropylene, Calcium Carbonate, PVC, Synthetic Rubber, Silicone, EVA NOT MADE WITH NATURAL RUBBER LATEX

### WARRANTY:

DJO, LLC will repair or replace all or part of the unit and its accessories for material or workmanship defects for a period of six months from the date of sale. To the extent the terms of this warranty are inconsistent with local regulations, the provisions of such local regulations will apply.

### RX ONLY. INTENDED FOR SINGLE PATIENT USE.


 DJO, LLC
 I
 A DJO
 Global Company

 T
 800.336.6569
 D
 760.727.1280
 F
 800.936.6569

 1430
 Decision Street
 I
 Vista, CA 92081-8553
 I
 U.S.A.

 DJOGlobal com
 DJOGlobal com
 DJOGlobal com
 DJOGlobal com
 DJOGlobal com







Individual results may vary. Neither DJO, LLC nor any of its subsidiaries dispense medical advice. The contents of this document do not constitute medical, legal, or any other type of professional advice. Rather, please consult your healthcare professional for information on the courses of treatment, if any, which may be appropriate for you.